



AVATAR
Same Day Surgery

Rapid-Recovery After Same Day Surgery
TOTAL HIP REPLACEMENT



Thomas V. King, MD
Moby Parsons, MD

THE
KNEE • HIP • SHOULDER
CENTER

333 Borthwick Ave, Suite 301
Portsmouth, New Hampshire 03801

(603) 431-5858



AVATAR *Same Day Surgery*

AVATAR[®] is a new approach that gives each person customized navigation through the complex process of total joint replacement. **AVATAR**[®] stands for the Alignment of Vital Assets To Accelerate Recovery. **AVATAR**[®] is a comprehensive care plan lead by a team of experienced specialists that guides patients through the process of rapid recovery and out-patient surgery. **AVATAR**[®] uses the best technology and surgical technique, selects the best professionals (surgical team, anesthetist, physical therapist), and ensures the best pain control to achieve faster recovery and a better outcome for life. **AVATAR**[®] has an exceptional track record of over 1000 patients who have done better with a personalized, customized total joint experience.

MY AVATAR[®] is the person you dream of being after your arthritis pain is gone. That person can walk, do stairs, sleep, play with pets or grandkids, play golf, ski, play tennis, hunt, fish and pursue and all activities without pain. That new person is your **AVATAR**[®].

HOW?

With new anesthetic techniques and minimally invasive surgery an overnight hospital stay is unnecessary for appropriately selected patients. The use of non-narcotic pain relievers, regional anesthesia (nerve blocks) and medication to limit blood loss make it possible for patients to be dressed, walking, doing stairs and heading home within a few hours after surgery.

WHO?

Healthy, motivated individuals who don't require a hospital stay to manage other health problems. After surgery, patients have to demonstrate that they are sufficiently recovered to be functional and safe at home. Patients are cleared for discharge to home by an experienced physical therapist, nursing staff and Surgeon. A total joint coach/ support person must be available to drive the patient home and spend the night after surgery

LESS PAIN

Pain is the major factor limiting rapid rehabilitation after total joint surgery. We cannot make surgery "pain-free" however by making surgery less invasive, by limiting the use of general anesthesia, and by enhancing our use of regional nerve blocks and longer-lasting local anesthetics (Exparel), rehabilitation and the return to normal activity can be accelerated. We still believe in the adage "no pain- no gain" and will encourage patients to push themselves maximally despite discomfort. Oral pain relievers are still necessary. An individualized postoperative medication schedule will be developed for each patient using multi-modal medications to limit the use of narcotics.

PREPARATION

A stepwise plan starts approximately 4 weeks before the date of surgery. Our office will facilitate this plan for you, this includes:

- Scheduling appropriate labwork, EKG and chest x-ray if needed
- Medical clearance with your Primary Care Physician
- Your PCP may want you to have further testing or consultations with other doctors.

PRE-HAB

Prior to surgery you may have an appointment with physical therapy for instruction on preparatory muscle and joint exercises. Following this exercise program will be your responsibility and will help you get ready for surgery and introduce you to your outpatient physical therapist.

HOME PHYSICAL THERAPY

A preoperative visit, or phone call from a home physical therapist will be ordered by our office. This is to help ensure a safe environment, orientation for family and other caregivers, rehearsal of the rehabilitation protocol at home. Our office will select the agency and send the referral.

ADAPTIVE OR DME

You should have a walker, crutches and a cane available. There are many community resources where these are available at little or no charge. Each community may have different resources, such as the Rye Lions Club, Kittery Lions Club, Care Pharmacy in Rochester, Keene Medical Portsmouth, Rite Aid pharmacies etc. *Please remember to bring your walker on the day of surgery.

ORTHOPEDIC PREOPERATIVE VISIT

You will have a pre-operative visit at your Surgeon's office approximately one week before surgery. This is to review the information from your appointment with your PCP and answer any questions.

- A medication schedule will be developed and reviewed with you.
- Postoperative prescriptions will be given to fill ahead of time.
- We recommend that you have Prilosec 20mg, Tylenol 500mg, enteric coated aspirin 325mg.
- You may want to have stool softeners and laxatives on hand.
- Let your surgeon's office know if you have motion sickness or have had difficulty with nausea after other surgery. They may want to prescribe something to help reduce the chance you will feel sick after your joint replacement.

PREOPERATIVE MEDICATION

- STOP taking anti inflammatory medications one week before surgery. *This includes Aleve, Advil, ibuprofen, Motrin, Voltaren and others.*
- You may continue taking aspirin if you are already on aspirin.
- If you have any questions call your Surgeon's office.

- START taking Tylenol 1000mg three times a day, beginning 3 days before your surgery date.
- DO NOT TAKE TYLENOL THE MORNING OF YOUR SURGERY.
- You will receive IV Tylenol on the day of your surgery.
- Starting 5 days prior to surgery, wash with **Hibiclens** when you shower (available over the counter at your pharmacy). This is to reduce the bacterial count on your skin.
- We recommend soaping up with Hibiclens in the shower, turn off the water for 2-3 minutes and then rinse. Avoid head and face.

SURGERY

One to two days prior to your surgery, you will receive a call from the Surgery Center or Hospital, letting you know your arrival time.

- No food after midnight the evening before your surgery.
- You may drink clear liquids such as water, Gatorade or apple juice until 3 hours before your arrival time.
- We encourage patients to hydrate for several days prior to surgery. This helps reduce feeling dehydrated after surgery and symptoms that may go along with that, such as dizziness, low blood pressure.
- Be mindful of your diet the week prior to surgery. Healthy foods including protein and good carbohydrates will help with energy, stamina and healing.

PRE-OP SUGGESTIONS

- Remove nail polish the day before surgery
- Consider making meals ahead of time and freezing
- Consider clean sheets on your bed for optimum hygiene
- Try to sleep well the night before surgery

THE DAY OF SURGERY

Upon arrival to the Surgery Center or Hospital there will be an evaluation by the nurses, the Anesthesiologist and your Surgeon.

- Once your IV is started you will have regional anesthesia (nerve blocks) and a spinal anesthetic
- You will receive preoperative IV antibiotics.
- Surgery is usually about 2 hours in length.
- Following surgery you will be in the PACU, (post anesthesia care unit) you may have something to eat and drink.
- Your vital signs will be monitored and pain managed.
- You will be evaluated for your ability to move safely while the nerve blocks are wearing off.
- You will work with an experienced physical therapist for safety transferring from sit to stand, safety with walking and practice stair climbing.
- Your vital signs will continue to be monitored.
- You will be assessed for your readiness for discharge to home if your plan was for same day surgery.
- Feeling weak immediately after surgery is normal, pace yourself, rest when you need to.

GOING HOME

HOME DAY 1

- Your total joint coach, (support person) will drive you home and be with you overnight. You can be up walking with assistance, going to the toilet, and doing stairs. Be safe, take it easy, pace yourself, remember you have just had surgery.
- You may move your operative leg gently to avoid stiffness.
- Stay hydrated, eat moderately if you feel well.
- Begin taking Aspirin 325mg every 12 hours daily with food, to prevent blood clots and continue for six weeks.
- Follow the medication schedule developed for you, to stay ahead of the pain.

HOME DAY 2

- Leave dressing in place, continue to pace yourself, stay hydrated and nourished.
- You will be seen by a home physical therapist who will help with safety on stairs and walking and work with you on range of motion, strength, and function.
- Continue with scheduled medications.

HOME DAY 3

- Home physical therapy
- Dressing change, incision may be open to air or covered with dressings provided. Incisions must be covered for showers for 10 days. Dr. Parsons' patients may have a dressing that remains in place for one week.
- No bathing/pool or hot tub until you have seen your surgeon 3 weeks after surgery. Your incision is closed with absorbable suture and steri strips. Steri strips will fall off over time.
- Call office for any incisional drainage or wound concerns.
- Continue scheduled medications/ avoid constipation

PAIN MEDICATIONS

(and other medications)

There are several different “classes” of pain receptors in your body. You will be prescribed non-narcotic medications that target these different receptors. By hitting more than one class of receptor the pain relief is enhanced. By following the multi-modal medication protocol/*medication schedule, that was developed for you, the medications work synergistically...more effectively.

Aleve 2 pills (inflammatory receptor) is taken twice daily with food.
Or ibuprofen including Advil or Motrin 2-3 tablets 2-3x/ daily with food

Tylenol 1000mg (“central” receptor) is taken 3 times per day (perhaps at 7am, 2pm and 9pm)

Tramadol 50mg (opioid receptor) is taken every 6 hours (perhaps at 7am, 2pm and 7pm and 12am) occasionally you can take an additional 50mg for “breakthrough” pain every 6 hours. Do not take more than 400mg of Tramadol in 24 hours. **some patients are not given RX for Tramadol due to contraindication with one of their regular medications

Aspirin 325mg is to prevent blood clots (not for pain) and should be taken twice daily for six weeks unless you are taking some other anticoagulant such as Coumadin, Eliquis, Plavix etc.

Oxycodone 5mg is a narcotic that is taken for more severe “breakthrough” pain if needed.

- You can take 5mg every 4-6 hours for moderate pain.
- 10mg every 4-6 hours for severe pain.

Zofran is commonly used to treat nausea and vomiting if this occurs.

Prilosec 20mg daily to protect your stomach while taking these medications.

You may need over the counter products to help avoid constipation.

FAQ'S

and post surgical recommendations

If you feel that you are faced with a life threatening emergency such as severe chest pain, severe shortness of breath, signs of symptoms of a heart attack or stroke call 911 and go directly to the emergency room. For all other emergencies call your Surgeon's office.

PAIN

Expect pain after total joint replacement. Pain may increase about 30 hours after surgery when the adductor canal nerve block wears off and then again in 2-3 days when the Exparel wears off. We recommend that you take scheduled medications as outlined to stay ahead of the pain. Call Dr. King's office with questions about pain.

NAUSEA

A common side effect of anesthesia and surgery. It may also be a side effect of constipation. It may also be a side effect of your medications.

SAFETY

Take care not to trip or fall. Move carefully and deliberately, use your walker, crutches or cane as instructed for at least the first three days. Use proper techniques on stairs to avoid falling. Adhere to any safe moving precautions you may have.

BREATHING

Keep your lungs expanded by deliberately taking deep breathes and coughing. If you were sent home from the hospital with an Incentive Spirometer continue using this regularly

FEVER

Temperatures up to and even above 101F is normal in the first few days after surgery. It does not require antibiotics. Call the office if you're experiencing fevers, sweats, or chills or have concerns.

DIET

You can consume a normal diet and are encouraged to eat foods that avert constipation.

CONSTIPATION

Is common with pain medications and must be avoided. Take a stool softener (such as Dulcolax) the day before surgery and continue throughout the first few days after surgery. Do not go for more than two days without a bowel movement. Laxatives are recommended, Metamucil, Milk of Magnesia and Miralax.

DIZZINESS and LIGHT-HEADEDNESS

Due to anesthesia, fluid loss, pain medications IS COMMON. Arising suddenly, rapid movements, and dehydration should be avoided. Initially move in stages and see how you feel, stay hydrated with water or electrolyte drinks such as Gatorade, Powerade, Pedialyte, these are readily available in grocery stores.

CLICKING

Total joint replacements are made from hard durable materials and you may feel or hear clicking with motion.

DRESSINGS

A bulky dressing can be removed on day 3. An island dressing, coverlet, or large Band-Aid can be applied after that before showering. It may be removed after showering.

SHOWERING AND BATHING

You can shower on day 3 with the incision covered after the bulky dressing has been removed. Remove the wet coverlet after showering and bathing. Do not immerse the incision in a bathtub or hot tub or pool for 3 weeks or until cleared by your surgeon's office.

DRIVING

You may drive when you feel safe and have discontinued any pain medications that can cause impairment. Your Physical Therapist may guide you regarding your readiness to drive.

FOR MORE INFORMATION

Office: Dr. King and Dr. Parsons	603-431-5858
Stratham Ambulatory Surgery Center	603-772-2076
Portsmouth Hospital	603-431-5110
SurgiCare (CPM Machines)	888-291-8901

PROVIDERS & OFFICE STAFF

Thomas V. King, MD

Moby Parsons, MD

Kathleen Leavitt, PA-C

Kimberly DeDeus, PA-C

Stacey Riley, PA-C

Dustin Forrest, APRN

Nancy La Liberte', Clinical Manager, Surgical Scheduler

Laurie Bentz, Registered Medical Assistant

Melanie Caruso, RT(R), Administrative Support

Jaime Griffin, Registered Medical Assistant

Chris Johnson, RT(R), Administrative Support

Izzy O'Hare, RT(R)

Wendy Rhoad, Senior Level Administrative Assistant/Receptionist



*avatar: the incarnation or embodiment
of an idea, concept or philosophy*



AVATAR
Same Day Surgery

333 Borthwick Ave, Suite 301
Portsmouth, New Hampshire 03801

(603) 431-5858